

MEDICAL STATEMENT FOR CHILDREN WITH LIFE-THREATENING ALLERGIES
To be Completed by Health Care Provider

Student's Name: _____ Birthdate: _____ Grade/Teacher _____

1. Please List LIFE-THREATENING allergens:

2. Please list foods to be omitted from diet, and/or substances to avoid:

3. Is this student trained, capable, and willing, and permitted to self-administer his/her Epinephrine Auto-Injector? yes no

4. Should this student sit at the nut-sensitive table during lunch? (Students who sit at the nut-sensitive table are asked to avoid bringing foods that contain peanuts and tree-nuts. Special attention is given to cleaning and sanitizing these tables.) yes no N/A

5. Will you allow an unlicensed person who is trained in Epinephrine Auto-Injector administration administer this child's Epinephrine Auto-Injector to him/her in the event of an emergency? yes no

6. I give permission to the School Nurse to share with appropriate school personnel information relative to my child's life-threatening allergy diagnosis and management? yes no

7. BFCCPS encourages the wearing of Medic-Alert or other similar type of jewelry, alerting others to the child's life-threatening allergy(ies).

Signature of Health Care Provider

Telephone Number

Date

Parent/Guardian Signature

Print Name

Date

**PLACE
PICTURE
HERE**

Name: _____ D.O.B.: _____

Allergy to: _____

Weight: _____ lbs. Asthma: [] Yes (higher risk for a severe reaction) [] No

NOTE: Do not depend on antihistamines or inhalers (bronchodilators) to treat a severe reaction. USE EPINEPHRINE.

Extremely reactive to the following foods: _____





THEREFORE:




[] If checked, give epinephrine immediately for ANY symptoms if the allergen was likely eaten.

[] If checked, give epinephrine immediately if the allergen was definitely eaten, even if no symptoms are noted.

FOR ANY OF THE FOLLOWING:

SEVERE SYMPTOMS





			
LUNG	HEART	THROAT	MOUTH
Short of breath, wheezing, repetitive cough	Pale, blue, faint, weak pulse, dizzy	Tight, hoarse, trouble breathing/ swallowing	Significant swelling of the tongue and/or lips

			OR A COMBINATION of symptoms from different body areas.
SKIN	GUT	OTHER	
Many hives over body, widespread redness	Repetitive vomiting, severe diarrhea	Feeling something bad is about to happen, anxiety, confusion	

↓ ↓ ↓

- INJECT EPINEPHRINE IMMEDIATELY.**
- Call 911.** Tell them the child is having anaphylaxis and may need epinephrine when they arrive.
 - Consider giving additional medications following epinephrine:
 - » Antihistamine
 - » Inhaler (bronchodilator) if wheezing
 - Lay the person flat, raise legs and keep warm. If breathing is difficult or they are vomiting, let them sit up or lie on their side.
 - If symptoms do not improve, or symptoms return, more doses of epinephrine can be given about 5 minutes or more after the last dose.
 - Alert emergency contacts.
 - Transport them to ER even if symptoms resolve. Person should remain in ER for at least 4 hours because symptoms may return.

MILD SYMPTOMS

			
NOSE	MOUTH	SKIN	GUT
Itchy/runny nose, sneezing	Itchy mouth	A few hives, mild itch	Mild nausea/ discomfort

FOR MILD SYMPTOMS FROM MORE THAN ONE SYSTEM AREA, GIVE EPINEPHRINE.

FOR MILD SYMPTOMS FROM A SINGLE SYSTEM AREA, FOLLOW THE DIRECTIONS BELOW:

- Antihistamines may be given, if ordered by a healthcare provider.
- Stay with the person; alert emergency contacts.
- Watch closely for changes. If symptoms worsen, give epinephrine.

MEDICATIONS/DOSES

Epinephrine Brand: _____

Epinephrine Dose: [] 0.15 mg IM [] 0.3 mg IM

Antihistamine Brand or Generic: _____


Antihistamine Dose: _____

Other (e.g., inhaler-bronchodilator if wheezing): _____

PARENT/GUARDIAN AUTHORIZATION SIGNATURE _____ DATE _____ PHYSICIAN/HCP AUTHORIZATION SIGNATURE _____ DATE _____


EPIPEN® (EPINEPHRINE) AUTO-INJECTOR DIRECTIONS

1. Remove the EpiPen Auto-Injector from the plastic carrying case.
2. Pull off the blue safety release cap.
3. Swing and firmly push orange tip against mid-outer thigh.
4. Hold for approximately 10 seconds.
5. Remove and massage the area for 10 seconds.



AUVI-Q™ (EPINEPHRINE INJECTION, USP) DIRECTIONS

1. Remove the outer case of Auvi-Q. This will automatically activate the voice instructions.
2. Pull off red safety guard.
3. Place black end against mid-outer thigh.
4. Press firmly and hold for 5 seconds.
5. Remove from thigh.



ADRENACLICK®/ADRENACLICK® GENERIC DIRECTIONS

1. Remove the outer case.
2. Remove grey caps labeled "1" and "2".
3. Place red rounded tip against mid-outer thigh.
4. Press down hard until needle penetrates.
5. Hold for 10 seconds. Remove from thigh.



OTHER DIRECTIONS/INFORMATION (may self-carry epinephrine, may self-administer epinephrine, etc.):

Treat the person before calling emergency contacts. The first signs of a reaction can be mild, but symptoms can get worse quickly.

EMERGENCY CONTACTS — CALL 911

RESCUE SQUAD: _____

DOCTOR: _____ PHONE: _____

PARENT/GUARDIAN: _____ PHONE: _____

OTHER EMERGENCY CONTACTS

NAME/RELATIONSHIP: _____

PHONE: _____

NAME/RELATIONSHIP: _____

PHONE: _____

PARENT/GUARDIAN AUTHORIZATION SIGNATURE _____

DATE _____